

June 22 1998
TAUP 2506-98

EVOLUTION EQUATIONS FOR HIGH PARTON DENSITY QCD

Eugene Levin

leving@post.tau.ac.il

School of Physics and Astronomy

Raymond and Beverly Sackler Faculty of Exact Science

Tel Aviv University, Tel Aviv, 69978, ISRAEL

Talk given at "Continuous Advances in QCD", Minnesota, April 16-19, 1998

Abstract. This talk is a brief report on our results in high parton density QCD. The main goal of this talk is to discuss the new evolution equation which was obtained and solved in Refs. [1] [2] and to share with you the physics can be recovered by this equation.

INTRODUCTION

In this talk, we briefly outline the main ideas, approaches and results which we developed in high parton density QCD (hdQCD). High parton density QCD involves a system of partons at short distances in which the QCD coupling constant α_S is small, but the density of partons becomes so large that we cannot apply the usual methods of perturbative QCD (pQCD). In essence, the theoretical problems here are non perturbative, but the origin of the non perturbative effects does not lie in long distances and large α_S , which are present in the confinement region.

First, we need to answer the question:

1. *Which parton density is high?*

The quantitative estimates of which density is high, we can obtain from the s -channel unitarity [3] which we can use in two different form:

1. $\sigma_{tot}(\gamma^*p) \leq \alpha_{em} 2\pi R^2$, where R is the size of the target, α_{em} is the fine structure constant (see Ref. [6] for explanation why we have α_{em}). This constraint can be rewritten as [3] [4] [5]

$$\kappa = \frac{3\pi\alpha_S x G(x, Q^2)}{Q^2 R^2} \leq 1. \quad (1)$$




FIGURE 1. Contours for $\langle \gamma \rangle$ and κ for the GRV'94 gluon density and HERA kinematic region.

2. $\sigma^{DD}(\gamma^*p) \leq \sigma_{tot}(\gamma^*p)$, where DD stands for diffractive dissociation. This inequality gives Eq. (1).

Therefore,

1. if κ is very small ($\kappa \ll 1$), we have a low density QCD in which the parton cascade can be perfectly described by the DGLAP evolution equations [7];

2. if $\kappa \leq 1$, we are in the transition region between low and high density QCD. In this region we can still use pQCD, but have to take into account the interaction between partons inside the partons cascade;

3. if $\kappa \geq 1$, we reach the region of high parton density QCD. which we are going to discuss here.

Before doing this we want to make a remark on parton densities in a nucleus. Taking into account that for nucleus $R_A = R_N \times A^{\frac{1}{3}}$ and $xG_A(x, Q^2) = AxG_N(x, Q^2)$, we can rewrite Eq. (1) as

$$\kappa_A = \frac{3\pi\alpha_S AxG_N(x, Q^2)}{Q^2 R_N^2 A^{\frac{2}{3}}} = A^{\frac{1}{3}} \kappa_N \leq 1. \quad (2)$$

Therefore, for the case of an interaction with nuclei, we can reach a hdQCD region at smaller parton density than in a nucleon (see Fig.1).

Fig.1 gives the kinematic plot (x, Q^2) with the line $\kappa = 1$, which shows that the hdQCD effect should be seen at HERA.

2. Two different theoretical approaches

To understand these two approaches we have to look at the picture of a high energy interaction in the parton model. In the parton approach the fast hadron decays into point-like particles (partons) long before (typical time $\tau \propto \frac{E}{\mu^2}$)




FIGURE 2. *The high energy interaction in the parton model.*

the interaction with the target. However, during this time τ , all partons are in the coherent state which can be described by means of a wave function. The interaction of the slowest (“wee”) parton with the target completely destroys the coherence of the partonic wave function.

The total cross section of such an interaction is equal to

$$\sigma_{tot} = N \times \sigma_0 \quad (3)$$

where

- $N = \text{flux (renormalized ?!) of “wee” partons ;}$
- $\sigma_0 = \text{the cross section of the interaction of one “wee” parton with the target .}$

One can see directly from Fig.2 that the number of “wee” partons is rather large and it is equal to

$$N \propto e^{\langle n \rangle} = \frac{1}{x^{\omega_0}} \text{ with } \langle n \rangle = \omega_0 \ln(1/x) ; \omega_0 = C\alpha_S . \quad (4)$$

We have to renormalize the flux of “wee” partons, since the total cross section is the number of interactions and if one has several “wee” partons with the same momenta they only give rise to one interaction.

If $N \approx 1$, we expect that the renormalization of the flux will be small, and we use an approach with the following typical ingredients:

- Parton Approach; • Shadowing Corrections; • Glauber Approach; • Reggeon-like Technique; • AGK cutting rules.

However, when $N \gg 1$, we have to change our approach completely from the parton cascade to one based on semiclassical field approach, since due to the uncertainty principle $\Delta N \Delta \phi \approx 1$, we can consider the phase as a small parameter. Therefore, in this kinematic region our magic words are:

- Semi-classical gluon fields; • Wiezsäcker-Williams approximation; • Effective Lagrangian for hdQCD; • Renormalization Wilson group Approach.

It is clear, that for $N \approx 1$ the most natural way is to approach the hdQCD looking for corrections to the perturbative parton cascade. In this approach the




FIGURE 3. The current situation in the battle for hdQCD.

pQCD evolution has been naturally included, and it aims to describe the transition region. The key problem is to penetrate into the hdQCD region where κ is large. Let us call this approach “pQCD motivated approach”.

For $N \gg 1$, the most natural way of doing is to use the effective Lagrangian approach, and remarkable progress has been achieved both in writing of the explicit form of this effective Lagrangian, and in understanding physics behind it [8]. The key problem for this approach was to find a correspondence with pQCD. This problem has been solved [9].

The **hottest news** of this conference is that these two approaches give the same evolution equation for the gluon structure function as reported by A. Kovner [10].

Fig.3 shows the current situation on the frontier line in the offensive on hdQCD.

3. Pluses and Minuses of pQCD motivated approach

The goal of this talk is to show you how we derive this evolution equation and explain what physics we can recover using this approach.

Let me list here the main advantages and shortcomings of this approach. I believe that the honest report on them will lead us to a deeper understanding of the main theoretical problems in hdQCD.

✚ -s:

1. Clear physical meaning of everything what we are doing;
2. Transparent parameters that we are using for a selection of the Feynman diagrams;
3. Obvious interrelation with usual DGLAP evolution and the Reggeon Calculus (“soft” phenomenology);

4. Understandable limitation of the approach which actually means that we guarantee the accuracy of our theoretical calculations. In other word, this approach has reached a stage of a normal theory with a regular procedure of how to calculate corrections with respect to small parameters.

5. It has a clear feedback for such interesting problems as higher twist contributions, violation of factorization and so on.

— :

There is **no hope** to obtain any formal (operator) approach to the problem and, therefore, to find a formal justification of the resummation technique. It means, that we can prove everything only in the framework of the Feynman diagrams and in the framework of the operation rules with them. However, it means that we ignore, at least partly, non perturbative effects and that we are not able to formulate a procedure of calculation of such effects.

Let me illustrate why it is very important an operator proof giving two examples.

1. In the pQCD motivated approach we heavily use the so called leading log $(1/x)$ approximation in pQCD (LL(x)A). The operation rule for LL(x)A is:

$$\begin{aligned} xG^{LL(x)A}(x, Q^2) &= \lim_{x \rightarrow 0} \sum_{n=1}^{\infty} \alpha_S^n M_n(x, Q^2) \\ &= \sum_{n=1}^{\infty} \alpha_S^n \lim_{x \rightarrow 0} M_n(x, Q^2) = \sum_{n=1}^{\infty} C_n(Q^2) (\alpha_S \ln(1/x))^n. \end{aligned} \quad (5)$$

We used the gluon structure function as an example. Eq. (5) shows that we changed the order of two operation : summation over n and limit at $x \rightarrow 0$. Do we have a proof for this? As far as I know, the answer is **no**. We consider as a strong argument in favor, the fact that the calculation in the next-to-leading order in LL(x)A gives a reasonable answer. The real proof has been done only recently [9] in the effective Lagrangian approach, which leads to the LL(x)A of pQCD.

2. One of the main non perturbative observable which enters any pQCD motivated calculation is so called the size of the target R (see Eq. (1)). We know the meaning of this parameter, namely, this is a correlation radius between two gluon inside a target at $x \approx 1$. In the effective Lagrangian approach we will be able to find an operator expression for this number, and to ask our friends from lattice QCD to calculate this value. I think this is the only reasonable way of how to take non perturbative QCD into account, at least, at present time.

NEW EQUATION

4. The GLR equation

The first attempt to take into account the parton - parton interaction in the pQCD motivated approach was done long ago [3]. It was based on the simple idea that there are two processes in a parton cascade (see Fig.2) : (i) the probability of the emission of an extra gluon is proportional to $\alpha_S \rho$ where ρ is the density of gluon in the transverse plane, namely

$$\rho = \frac{xG(x, Q^2)}{\pi R^2}; \quad (6)$$

and (ii) the annihilation of a gluon, or in other words a process in which the probability is proportional to $\rho^2 \times \sigma_{annihilation}$. $\sigma_{annihilation}$ can be estimated as $\sigma_{annihilation} \propto \alpha_S r^2$, where r is the size of the parton produced in the annihilation process. For deep inelastic scattering, $r^2 \propto \frac{1}{Q^2}$. Therefore, in the parton cascade we have

$$Emission (1 \rightarrow 2) : probability = P^{emission} \propto \alpha_S \rho ; \quad (7)$$

$$Annihilation (2 \rightarrow 1) : probability = P^{annihilation} \propto \alpha_S^2 \frac{1}{Q^2} \rho^2 . \quad (8)$$

At $x \sim 1$ only emission of new partons is essential, because $\rho \ll 1$ and this emission is described by the DGLAP evolution equations. However, at $x \rightarrow 0$ the value of ρ becomes so large that the annihilation of partons becomes important, and so the value of ρ is diminished. The competition of these two processes we can write as an equation for the number of partons in a phase space cell¹ ($\Delta y = \Delta \ln(1/x) \Delta \ln(Q^2/Q_0^2)$):

$$\frac{\partial^2 \rho}{\partial \ln(1/x) \partial \ln(Q^2/Q_0^2)} = \frac{\alpha_S N_c}{\pi} \rho - \frac{\alpha_S^2 \tilde{\gamma}}{Q^2} \rho^2 , \quad (9)$$

or in terms of the gluon structure function $xG(x, Q^2)$

$$\frac{\partial^2 xG(x, Q^2)}{\partial \ln(1/x) \partial \ln(Q^2/Q_0^2)} = \frac{\alpha_S N_c}{\pi} xG(x, Q^2) - \frac{\alpha_S^2 \tilde{\gamma}}{\pi R^2 Q^2} (xG(x, Q^2))^2 . \quad (10)$$

This is the GLR equation which gave the first theoretical basis for the consideration of hdQCD. This equation describes the transition region at very large values of Q^2 , but a glance at Fig.1 shows that we need a tool to penetrate the kinematic region of moderate and even small Q^2 .

5. Glauber - Mueller Approach

We found that it is very instructive to start with the Glauber approach to SC. The idea of how to write the Glauber formula in QCD was originally formulated in two papers Ref. [11] and Ref. [5]. However, the key paper for our problem is the second paper of A. Mueller, who considered the Glauber approach for the gluon structure function. The key observation is that the fraction of energy, and the transverse coordinates of the fast partons can be considered as frozen, during the high energy interaction with the target [11] [5]. Therefore, the cross section of the absorption of gluon(G^*) with virtuality Q^2 and Bjorken x can be written in the form:

$$\sigma_{tot}^N(G^*) = \quad (11)$$

$$\int_0^1 dz \int \frac{d^2 r_t}{2\pi} \int \frac{d^2 b_t}{2\pi} \Psi_{\perp}^{G^*}(Q^2, r_t, x, z) \sigma_N(x, r_t^2) [\Psi_{\perp}^{G^*}(Q^2, r_t, x, z)]^* ,$$

where z is the fraction of energy which is carried by the gluon, $\Psi_{\perp}^{G^*}$ is the wave function of the transverse polarized gluon, and $\sigma_N(x, r_t^2)$ is the cross section of the

¹⁾ We will argue why we chose this cell below in section 6.

interaction of the GG - pair with transverse separation r_t with the nucleus. Mueller showed that Eq. (11) can be reduced to

$$xG^{MF}(x, Q^2) = \frac{4}{\pi^2} \int_x^1 \frac{dx'}{x'} \int_{\frac{1}{Q^2}}^\infty \frac{d^2 r_t}{\pi r_t^4} \int_0^\infty \frac{d^2 b_t}{\pi} 2 \left\{ 1 - e^{-\frac{1}{2} \sigma_N^{GG}(x', r_t^2) S(b_t^2)} \right\} \quad (12)$$

with

$$\sigma_N^{GG} = \frac{4 \pi^2 \alpha_S}{3} r_t^2 xG^{DGLAP}\left(x, \frac{1}{r_t^2}\right)$$

and profile function $S(b_t)$ chosen in the Gaussian form: $S(b_t^2) = \frac{1}{\pi R^2} e^{-\frac{b_t^2}{R^2}}$.

Obviously, the Mueller formula has a defect, namely, only the fastest partons (GG pairs) interact with the target. This assumption is an artifact of the Glauber approach, which looks strange in the parton picture of the interaction. Indeed, in the parton model we expect that all partons not only the fastest ones should interact with the target. At first sight we can solve this problem by iteration Eq. (12) (see Ref. [1]). It means that the first iteration will take into account that not only the fastest parton, but the next one will interact with the target, and so on.

6. Why equation?

We would like to suggest a new approach based on the evolution equation to sum all SC. However, we want to argue first, why an equation is better than any iteration procedure. To illustrate this point of view, let us differentiate the Mueller formula with respect to $y = \ln(1/x)$ and $\xi = \ln Q^2$. It is easy to see that this derivative is equal to

$$\frac{\partial^2 xG(x, Q^2)}{\partial y \partial \xi} = \frac{4}{\pi^2} \int db_t^2 \left\{ 1 - e^{-\frac{1}{2} \sigma(x, r_\perp^2 = \frac{1}{Q^2}) S(b_t^2)} \right\}. \quad (13)$$

The advantages of Eq. (13) are

- Everything enters at small distances ;
- Everything is under theoretical control ;
- Everything that is not known (mostly non perturbative) is hidden in the initial and boundary condition.

Of course, we cannot get rid of our problems changing the procedure of solution. Indeed, the non-perturbative effects coming from the large distances are still important, but they are absorbed in the boundary and initial conditions of the equation. Therefore, an equation is a good (correct) way to separate what we know (small distance contribution) from what we don't (large distance contribution).

7. Equation

We suggest the following way to take into account the interaction of all partons in a parton cascade with the target. Let us differentiate the b_t -integrated Mueller formula of Eq. (12) in $y = \ln(1/x)$ and $\xi = \ln(Q^2/Q_0^2)$. It gives

$$\frac{\partial^2 xG(y, \xi)}{\partial y \partial \xi} = \frac{2Q^2 R^2}{\pi^2} \left\{ C + \ln(\kappa_G(x', Q^2)) + E_1(\kappa_G(x', Q^2)) \right\} , \quad (14)$$

where $\kappa_G(x, Q^2)$ is given by

$$\kappa(x, Q^2) = \frac{N_c \alpha_S \pi}{2Q^2 R^2} xG(x, Q^2) . \quad (15)$$

Therefore, we consider κ_G on the l.h.s. of Eq. (14) as the observable which is written through the solution of Eq. (14).

Eq. (14) can be rewritten in the form (for fixed α_S)

$$\frac{\partial^2 \kappa_G(y, \xi)}{\partial y \partial \xi} + \frac{\partial \kappa_G(y, \xi)}{\partial y} = \frac{N_c \alpha_S}{\pi} \left\{ C + \ln(\kappa_G) + E_1(\kappa_G) \right\} \equiv F(\kappa_G) . \quad (16)$$

This is the equation which we propose.

8. Nice properties of the equation:

This equation has the following desirable properties:

1. It sums all contributions of the order $(\alpha_S y \xi)^n$, absorbing them in $xG(y, \xi)$, as well as all contributions of the order of κ^n . Therefore, this equation solves the old problem, formulated in Ref. [3], and for $N_c \rightarrow \infty$ Eq. (16) gives the complete solution to our problem, summing all SC ;
2. The solution of this equation matches with the solution of the DGLAP evolution equation in the DLA of perturbative QCD at $\kappa \rightarrow 0$;
3. At small values of κ ($\kappa < 1$) Eq. (16) gives the GLR equation. Indeed, for small κ we can expand the r.h.s of Eq. (16) keeping only the second term. Rewriting the equation through the gluon structure function we obtain the GLR equation [3] with the coefficient in front of the second term as calculated by Mueller and Qiu [4] ;
4. The first iteration of this equation gives the Mueller formula (see Ref. [5]) ;
5. In general, everything that we know about SC is included in Eq. (16) ;
6. As has been mentioned, Eq. (16) was reproduced by the effective Lagrangian approach (see A. Kovner talk at this conference [10] .

9. The theory status of the equation

Here, we list all small (large) parameters that have been used to obtain Eq. (16):

$$\alpha_S \ln(1/x) \ln(Q^2/Q_0^2) \approx 1 ; \quad (17)$$

$$\alpha_S \ln(1/x) < 1 ; \quad \alpha_S \ln(Q^2/Q_0^2) < 1 ; \quad (18)$$

$$\alpha_S \ll 1 . \quad (19)$$

- These equations (Eq. (17) - Eq. (19)) give the conditions when we can absorb all $(\alpha_S \ln(1/x) \ln(Q^2/Q_0^2))^n$ - terms in the gluon structure function both in Eq. (16) and in Eq. (12).

$$\kappa \leq \frac{1}{\alpha_S} . \quad (20)$$

- This is the main limitation of our resummation which stems from the contribution of so called enhanced diagrams (see Ref. [1] for detail).

$$N_c \gg 1 . \quad (21)$$

- where N_c is the number of colours. We follow the Veneziano principle [12] of large N_c resummation: we select the diagrams in leading order with respect to $(\alpha_s N_c)^n$ in each topologic configuration (see Ref. [1] and references therein for a detail discussion of this point).

SOLUTION

10. Asymptotic solution

A first observation is the fact that Eq. (16) has a solution which depends only on y . Indeed, one can check that $\kappa = \kappa_{asympt}(y)$ is the solution of the following equation

$$\frac{d\kappa_{asympt}}{dy} = F(\kappa_{asympt}) . \quad (22)$$

The solution to the above equation is

$$\int_{\kappa_{asympt}(y=y_0)}^{\kappa_{asympt}(y)} \frac{d\kappa'}{F(\kappa')} = y - y_0 . \quad (23)$$

one can see that Eq. (23) leads to

$$\begin{aligned} \kappa_{asympt}(y) &\rightarrow \alpha_S y \ln(\alpha_S y) \text{ at } \alpha_S y \gg 1 ; \\ \text{or } xG(x, Q^2) &\propto Q^2 R^2 \ln(1/x) \ln(\alpha_S \ln(1/x)) \text{ at } x \rightarrow 0 \end{aligned} \quad (24)$$

We claim this solution is the asymptotic solution of Eq. (16). To prove this we have to consider the stability of the asymptotic solution. It means, that we solve our general equation looking for the solution of the form

$$\kappa = \kappa_{asympt}(y) + \Delta\kappa(y, \xi - \xi_0) , \quad (25)$$

where $\Delta\kappa$ is small ($\Delta\kappa \ll \kappa_{asympt}$) but an arbitrary function at $\xi = \xi_0$. We have to prove that Eq. (25) will not lead to big $\Delta\kappa$ ($\Delta\kappa \gg \kappa_{asympt}$) at large ξ . The following linear equation can be written for $\Delta\kappa$

$$\frac{\partial^2 \Delta\kappa(y, \xi)}{\partial y \partial \xi} + \frac{\partial \Delta\kappa(y, \xi)}{\partial y} = \frac{dF(\kappa)}{d\kappa} \Big|_{\kappa=\kappa_{asympt}(y)} \Delta\kappa(y, \xi) . \quad (26)$$

In Ref. [1] it was proven, that the solution of Eq. (25) is much smaller than κ , since $\frac{dF(\kappa)}{d\kappa} \rightarrow 0$ at large y .

Therefore, the asymptotic solution has a chance to be the solution of Eq. (16) in the region of very small x . To prove this, we need to solve the equation in the wide kinematic region starting with the initial condition. We managed to do this only in the semiclassical approach.

11. The semiclassical approach

In the semiclassical approach the solution of Eq. (16) is supposed to be of the form

$$\kappa = e^S, \quad (27)$$

where S is a function with partial derivatives: $\frac{\partial S}{\partial y} = \omega$ and $\frac{\partial S}{\partial \xi} = \gamma$ which are smooth functions of y and ξ .

For an equation in the form $F(\xi, y, S, \gamma, \omega) = 0$, we can introduce the set of characteristic lines $(\xi(y), S(y), \omega(y), \gamma(y))$, which satisfy a set of well defined equations (see, for example, Refs. [13], [14] for the method). In Ref. [1] we developed a detailed calculation of the solution. The set of equations is:

$$\frac{d\xi}{dy} = \frac{\Phi(S)}{(\gamma + 1)^2}; \quad \frac{dS}{dy} = \frac{2\gamma + 1}{(\gamma + 1)^2}\Phi(S); \quad \frac{d\gamma}{dy} = \Phi'S \frac{\gamma}{\gamma + 1}; \quad (28)$$

where $\Phi'_S = \frac{\partial \Phi(S)}{\partial S}$. The initial conditions for this set of equations

$$S_0 = \ln \kappa_{in}(y_0, \xi_0); \quad \gamma_0 = \left. \frac{\partial \ln \kappa_{in}(y_0, \xi)}{\partial \xi} \right|_{\xi=\xi_0}. \quad (29)$$

The main properties of these equations have been considered in Ref. [1] analytically. Here, however, we restrict ourselves mostly to the numerical solution of these equations.

Fig. 4 shows the behaviour of trajectories and γ value for the solution. One can see that the trajectories can be divided in two groups: (i) first with $\gamma < -\frac{1}{2}$, and all of them approach the trajectories of the DGLAP evolution equation at large Q^2 ; and (ii) second one with $\gamma > -\frac{1}{2}$ and the solution of Eq. (16) on these trajectories tends to the asymptotic solution at $y \gg 1$.

We consider this figure as a nice illustration of our statement that the high energy (low x) limit of the general solution to Eq. (16) is the asymptotic solution.

12. Comparison with other approaches

In Fig.5 we plot our calculation for the DGLAP evolution equation, for the Glauber - Mueller approach, for the GLR equation and for the new evolution equation. We can conclude, that:

1. SC even in the Glauber - Mueller approach are essential for the gluon structure function ;




FIGURE 4. The trajectories (a) and the γ values (b) for the solution of the generalized evolution equation for nucleon with $R^2 = 5 \text{ GeV}^{-2}$.




FIGURE 5. The current situation in the battle for $hdQCD$.

2. The Glauber - Mueller approach considerably underestimates the value of SC ;
3. The GLR equation leads to stronger SC than the solution to Eq. (16) ;
4. The new evolution equation does not reproduce the saturation of the gluon density in the region of small x which the GLR equation leads to .

SUMMARY

13. Where are SC?

We firmly believe that the new evolution equation gives the correct way of evaluation of the value of SC. However, the difficult question arises: why the SC have not been seen at HERA? Our answer is:

1. The value of SC is rather large but only in the gluon structure function while their contribution to F_2 is rather small [1], and cannot be seen on the background of the experimental errors ;

2. The theoretical determination of the gluon structure function is not very precise and we evaluate the errors as 50% [15] and the SC in xG is hidden in such large errors ;

3. The statement that SC have not been seen is also not quite correct. Our estimates [16] show that the contribution of SC is rather large in the F_2 slope and incorporating the SC one can describe the recent experimental data [17] on $\frac{\partial F_2(x, Q^2)}{\partial \ln(Q^2/Q_0^2)}$ (see Ref. [16] for detail).

14. Last words

I hope that I gave you a brief but to some extent full review of our approach. Obviously, you can obtain more information from two large papers of Ref. [1]. Deeply in my heart, I firmly believe that there will be a bright future for the effective Lagrangian approach in which one combines the physics of hdQCD, with the formal methods of the quantum field theory, and gives a way to incorporate the lattice calculation for the non-perturbative observables. However, the parameter or better to say a new scale $Q_0^2(x)$ which is the solution of the equation $\kappa(x, Q_0^2(x)) = 1$ only can be found in the pQCD motivated approach, since the effective Lagrangian was only derived and justified assuming this new scale. The fact that the correct equation was first found in this approach is also not accidental . since the pQCD motivated approach has a great advantage : clear understanding of physics of hdQCD.

REFERENCES

1. A.L. Ayala, M.B. Gay Ducati and E.M. Levin: *Nucl. Phys.* **B493**(1997) 305; **B510** (1998) 355.
2. E. Gotsman, E. Levin and U. Maor: *Phys. Lett* **B379** (1996) 186; **B430** (1997) 120; *Nucl. Phys.* **B464** (1996) 251; **B493** (1997) 354; hep-ph/9712517.

3. L. V. Gribov, E. M. Levin and M. G. Ryskin: *Phys.Rep.* **100** (1983) 1.
4. A.H. Mueller and J. Qiu: *Nucl. Phys.* **B268** (1986) 427.
5. A.H.Mueller: *Nucl. Phys.* **B335** (1990) 115.
6. V.N. Gribov: *Sov. Phys. JETP* **30** (1970) 709.
7. V.N. Gribov and L.N. Lipatov: *Sov. J. Nucl. Phys.* **15** (1972) 438; L.N. Lipatov: *Yad. Fiz.* **20** (1974) 181; G. Altarelli and G. Parisi: *Nucl. Phys.* **B126** (1977) 298; Yu.L. Dokshitzer: *Sov. Phys. JETP* **46** (1977) 641.
8. L. McLerran and R. Venugopalan: *Phys. Rev.* **D49** (1994) 2233,3352, **D50** (1994) 2225, **D53** (1996) 458; J. Jalilian-Marian, A. Kovner, A. Leonidov and H. Weigert: hep - ph/9701284 hep-ph/9706377; J. Jalilian-Marian, A. Kovner, L. McLerran and H. Weigert: *Phys. Rev.* **D55** (1997) 5414; Yu. Kovchegov: *Phys. Rev.* **D54** (1996) 5463; Yu. Kovchegov and A.H. Mueller: hep-ph/9802440.
9. J. Jalilian-Marian, A. Kovner, A. Leonidov and H. Weigert: hep - ph/9701284 hep-ph/9706377.
10. A. Kovner: *these proceedings*.
11. E.M. Levin and M.G. Ryskin: *Sov. J. Nucl. Phys.* **45** (1987) 150.
12. G.Veneziano: *Phys. Lett.* **B52**(1974) 220; *Nucl. Phys.* **B94** (1974) 365; G. Marchesini and G. Veneziano: *Phys. Lett.* **B56**(1975) 271; M. Ciafaloni, G.Marchesini and G. Veneziano: *Nucl. Phys.* **B98** (1975) 472.
13. J.C. Collins and J. Kwiecinski: *Nucl. Phys.* **B335** (1990) 89.
14. J. Bartels, J. Blumlein and G. Shuler: *Z. Phys.* **C50** (1991) 91.
15. E. Levin: TAUP 2450-97, DESY 97-171, hep-ph/9709226.
16. E. Gotsman, E. Levin and U. Maor: TAUP 2471/97, hep-ph/9712517.
17. A. Caldwell: Invited talk in the DESY Theory Workshop, DESY, October 1997.